

SUMMARY OF PRODUCTS CHARACTERISTICS. 1. NAME OF THE MEDICINAL PRODUCT: FOLIC ACID 1. Qualitative and quantitative: Each tablet contains: Folic acid (INN), 400 micrograms; cyanocobalamin (INN) (vitamin B12), 2 micrograms, excipients, see List of excipients, qs

2. Pharmaceutical form: Yellow round tablets.

3. Clinical data

3.1. Indications therapeutic: Prevention of deficiency states of folate and vitamin B12 (cyanocobalamin)_in women of childbearing age who have planned a pregnancy for one month before conception and three months thereafter, as prophylaxis against neural tube defects.

3.2. Dosage We recommend the following dosing regimen: one tablet daily, administered before meals

3.3 Contraindications. There are no known specific contraindications dosage.

3.4 Special warnings and precautions. There is no warning or caution in their use worth highlighting. This product contains 23 mg of lactose per tablet which should be considered in patients with absorption of glucose or galactose, galactosaemia or lactase deficiency, although the amount present in the preparation is probably not enough to cause symptoms of intolerance.

3.5 Interactions. Folidoce not be administered together with methotrexate, as this acts as a folic acid antagonist by inhibiting the enzyme hidrofolatoreductasa. The combination of antiepileptic Folidoce with the group of hydantoins may decrease the effects of the latter.

3.6 Pregnancy and lactation Folidoce is indicated for the prevention of deficiencies_of vitamin B12 and folic acid before and during pregnancy. However, if you are pregnant or think you might be, consult your doctor.

3.7 Effects on ability to drive: No results.

3.8 Undesirable effects

3.9 Overdose. Very high doses of folic acid and vitamin B12 to those Folidoce is well tolerated, so it is not even afraid of poisoning by accidental massive ingestion.

4. PHARMACEUTICAL

4.1 List of excipients Type A starch sodium, microcrystalline cellulose, magnesium stearate, lactose monohydrate.

4.2 Incompatibilities. It presents the dose of administration.

4.3 Period of validity. Two years.

4.4 Storage Keep cool and dry.

4.5 Pack Contents Carton containing 28 tablets in blister white PVC / aluminum.